

Strange form factors in the context of SAMPLE, HAPPEX, and A4 experiments

António Silva^(1,2) *, Hyun-Chul Kim⁽³⁾ †, and Klaus Goeke⁽¹⁾ ‡

(1) *Institut für Theoretische Physik II,
Ruhr-Universität Bochum,
D-44780 Bochum, Germany*

(2) *Departamento de Física and Centro de Física Computacional,
Universidade de Coimbra,
P-3000 Coimbra, Portugal*

(3) *Department of Physics,
Pusan National University,
609-735 Pusan, Republic of Korea*

(Dated: September 2001)

Abstract

The strange properties of the nucleon are investigated within the framework of the SU(3) chiral quark-soliton model assuming isospin symmetry and applying the symmetry-conserving SU(3) quantization. We present the form factors $G_{E,M}^0(Q^2)$, $G_M^Z(Q^2)$ and the electric and magnetic strange form factors $G_{E,M}^s(Q^2)$ incorporating pion and kaon asymptotics. The results show a fairly good agreement with the recent experimental data from the SAMPLE and HAPPEX collaborations. We also present predictions for future measurements including the A4 experiment at MAMI (Mainz).

PACS numbers: 12.40.-y, 14.20.Dh

Keywords : strange vector form factors; chiral quark-soliton model

* E-mail address: Antonio.Silva@tp2.ruhr-uni-bochum.de

† E-mail address: hchkim@pnu.edu

‡ E-mail address: Klaus.Goeke@tp2.ruhr-uni-bochum.de

1. The understanding of the role that the sea quarks play in the properties of the nucleon and other hadrons is of fundamental importance to the description of their internal dynamics. With this respect, the determination of the strange contribution to the properties of the nucleon, for instance to its magnetic moment, charge and magnetic radii, is very important since there are no valence strange quarks in the nucleon. The aforementioned contributions to the properties of the nucleon are all included in its strange form factors, which means that the question may be rephrased as to whether or not the strange form factors contribute to e.g. the nucleon electromagnetic form factors. Since it became clear that such question could be addressed in weak neutral current experiments [1, 2] a great deal of effort both theoretical and experimental has been put into this aspect of the strange content of the nucleon.

On the experimental side examples of the ongoing efforts focusing on the strange vector form factors are the measurements by the SAMPLE [3, 4] and HAPPEX [5] collaborations. The most recent measurement by the SAMPLE collaboration [4] for the strange magnetic form factor finds (Q^2 in $(\text{GeV}/c)^2$)

$$G_M^s(Q^2 = 0.1) = (+0.14 \pm 0.29 \text{ (stat.)} \pm 0.31 \text{ (syst.)}) \text{ n.m.} \quad (1)$$

It is obtained from knowledge both on the neutral weak magnetic form factor G_M^Z measured in parity-violating elastic electron-proton scattering and on the electromagnetic form factors $G_M^{p\gamma}$, $G_M^{n\gamma}$ using the relation

$$G_M^Z = \left(1 - 4 \sin^2 \theta_W\right) G_M^{p\gamma} - G_M^{n\gamma} - G_M^s, \quad (2)$$

where θ_W is the weak mixing angle determined experimentally [6]: $\sin^2 \theta_W = 0.23147$. The HAPPEX collaboration also announced results on the measurement of the strange vector form factors [5]: From the parity-violating polarized electron scattering asymmetry A_{th} they arrive after various approximations at

$$\frac{(G_E^0 + 0.392G_M^0)}{(G_M^{p\gamma}/\mu_p)}(Q^2 = 0.477) = 1.527 \pm 0.048 \pm 0.027 \pm 0.011. \quad (3)$$

The first error is statistical, the second is systematic and the last comes from the uncertainty of the proton axial form factor [5]. The available data for the electromagnetic form factors have access to the strange vector form factors through

$$G_{E,M}^s = G_{E,M}^0 - G_{E,M}^{p\gamma} - G_{E,M}^{n\gamma} \quad (4)$$

with the result

$$(G_E^s + 0.392G_M^s)(Q^2 = 0.477) = 0.025 \pm 0.020 \pm 0.014. \quad (5)$$

Here, the first error is due to the errors in G_0 combined in quadrature and the second one arises from the electromagnetic form factors. Both procedures (2,4) show how uncertainties in the knowledge of the electromagnetic form factors propagate into the error bars of the results (1,5) for the strange form factors. In particular, the HAPPEX results show a much smaller error bar at the level of (3), which is extracted directly from the asymmetry A_{th} , than (5), obtained from Eq. (3) using Eq. (4). Unfortunately in the chiral quark-soliton model (χ QSM) we are presently not yet able to calculate the axial-vector form factor to

the same technical level as the electromagnetic ones. Thus we cannot calculate A_{th} but will focus on the expressions (3) and (5) for HAPPEX.

On the theoretical side there have been many attempts to predict the strange form factors, differing by the methods used to study the non-perturbative physics behind them: dispersion relations [7, 8, 9], Skyrme model [10], vector dominance models with $\omega - \phi$ mixing [11], Nambu–Jona-Lasinio soliton model [12], chiral bag model [13], meson exchange model [14], heavy baryon chiral perturbation theory [15, 16] and a chiral quark model [17]. The different physics going into these approaches explains the wide range of predictions obtained. Only the experimental data expected in the near future will allow to clarify the role of the mechanisms underlying these approaches. The non-perturbative lattice QCD calculations point towards a negative strange magnetic moment [18], but are not yet conclusive regarding the exclusion of positive values [19]. The strange form factors have already been investigated within the framework of the χ QSM [20]. However, it was later found that the collective SU(3) quantization used there for the χ QSM did not fulfil the Gell-Mann–Nishijima relation exactly, a fact which is caused by the inherent nonlocality in time of the collective operators in the χ QSM and the asymmetric transitions between the solitonic states and vacuum ones. We use in this study a procedure known as the symmetry-conserving quantization [21] for which the above problem does not arise. This new quantization has implications mainly at the level of the magnetic moments and hence is relevant for the present study.

2. The real Dirac and Pauli vector form factors are defined from the matrix elements of the quark current J_μ . In standard notation

$$\langle N(p') | J_\mu | N(p) \rangle = \bar{u}_N(p') \left[\gamma_\mu F_1(Q^2) + i\sigma_{\mu\nu} \frac{q^\nu}{2M_N} F_2(Q^2) \right] u_N(p), \quad (6)$$

where M_N and $u_N(p)$ are the nucleon mass and the nucleon spinor, respectively, and $Q^2 = -q^2 > 0$ is the square of the space-like four-momentum transfer, $q = p' - p$. In the present work we are interested in the form factors for the strange and baryonic currents. The strange quark current

$$J_\mu^s = \bar{s}\gamma_\mu s = J_\mu^B - J_\mu^Y \quad (7)$$

is expressed in terms of the baryonic J_μ^B and the hypercharge J_μ^Y currents:

$$J_\mu^B = \frac{1}{N_c} \bar{q}\gamma_\mu q, \quad (8)$$

$$J_\mu^Y = \frac{1}{\sqrt{3}} \bar{q}\gamma_\mu \lambda_8 q. \quad (9)$$

We prefer, however, to work with the electric and magnetic Sachs form factors, related to the previous ones by $G_E = F_1 - \tau F_2$, with $\tau = Q^2/(4M_N^2)$, and $G_M = F_1 + F_2$. Such a choice is more natural when working in the Breit frame and in the non-relativistic limit, as is our case, because the form factors are then expressed in a simple way by the time and space components of the current

$$\langle N'(J'_3)p' | J^0(0) | N(J_3)p \rangle = G_E(Q^2) \delta_{J'_3, J_3}, \quad (10)$$

$$\langle N'(J'_3)p' | J^k(0) | N(J_3)p \rangle = \frac{i}{2M_N} \epsilon^{klm} (\tau^l)_{J'_3 J_3} q^m G_M(Q^2). \quad (11)$$

The matrix element $\langle N'(J'_3)p' | J^\mu(0) | N(J_3)p \rangle$ is conserved in the χ QSM for the calculation of form factors described below. The form factors $G_{E,M}^0$ and $G_{E,M}^s$ here presented refer to the currents J_μ^B (8) and J_μ^s (7), respectively.

3. The starting point for the χ QSM [22] (for more details see [23, 24]) is the low-energy partition function in Euclidean space given by the functional integral over pseudo-Goldstone meson (π^a) and quark fields (ψ)

$$\mathcal{Z} = \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{D}\pi \exp \left[- \int d^4x \psi_f^\dagger D(\pi)_{fg} \psi_g \right] = \int \mathcal{D}\pi \exp(-S_{\text{eff}}[\pi]), \quad (12)$$

where $S_{\text{eff}}[\pi]$ denotes the effective action

$$S_{\text{eff}}[\pi] = -N_c \text{Tr} \ln D(\pi). \quad (13)$$

In these expressions Tr stands for the functional trace, N_c the number of colors, and D the Dirac differential operator in Euclidean space

$$D = \gamma_4 \left(\not{\partial} - \hat{m} - MU\gamma^5 \right) = \partial_4 + h(U) + m_8 \gamma_4 \lambda^8. \quad (14)$$

The λ^a are the Gell-Mann matrices normalized as $\text{tr}(\lambda^a \lambda^b) = 2\delta^{ab}$ and $\hat{m}$ is the matrix of the current quark mass given by $\hat{m} = \text{diag}(\bar{m}, \bar{m}, m_s) = m_1 \mathbf{1} + m_8 \lambda^8$ with $m_1 = (2\bar{m} + m_s)/3$ and $m_8 = (\bar{m} - m_s)/\sqrt{3}$ assuming isospin symmetry, *i.e.* with $\bar{m} = m_u = m_d$. The constituent quark mass M arises from the spontaneous breakdown of chiral symmetry and is considered to be constant and the only free parameter of the model. The proper-time regularization is chosen in order to complete the definition of the model. For U we assume a structure corresponding to Witten's so-called trivial embedding of $\text{SU}(2)$ into $\text{SU}(3)$:

$$U_{\text{SU}(3)} = \begin{pmatrix} U_{\text{SU}(2)} & 0 \\ 0 & 1 \end{pmatrix}, \quad (15)$$

with the $\text{SU}(2)$ hedgehog given by

$$U_{\text{SU}(2)} = \exp[i\gamma_5 \mathbf{n} \cdot \boldsymbol{\tau} P(r)]. \quad (16)$$

The partition function $\mathcal{Z}$ (12) can not be treated exactly as far as the integration over the pseudoscalar fields is concerned and thus requires some approximation framework, chosen in this model to be that of a large number of colors N_c . In the large N_c limit the Euclidean partition function $\mathcal{Z}$ can be evaluated in the saddle-point approximation, which corresponds at the classical level to finding the profile function $P(r)$ in Eq. (16) which makes the action stationary. This means in practice the numerical solution of the functional fixed point equation resulting from $\delta S_{\text{eff}}/\delta P(r) = 0$. Such procedure yields a classical field U_c built from a set of single quark energies and corresponding states, E_n and Ψ_n , pertaining to the one-particle eigenvalue problem for the Hamiltonian $h(U)$ in Eq. (14). This mean-field object does not have the quantum numbers of the nucleon state, however. To obtain the nucleon quantum numbers it is necessary to extract them from U_c by the semiclassical quantization of the rotational and translational zero-modes: The zero modes are treated exactly within the path integral formalism by introducing collective coordinates (see [23] for further details.) Explicitly, the rotational and translational zero modes are taken into account by means of the replacement

$$U(\mathbf{x}, t) = R(t) U_c(\mathbf{x} - \mathbf{Z}(t)) R^\dagger(t), \quad (17)$$




FIG. 1: The form factors G_E^0 and G_M^0 (in physical n.m.) as a function of Q^2 . The solid curve and dashed curves represent the results for the kaon ($\mu = 490$ MeV) and pion ($\mu = 140$ MeV) asymptotic tails, respectively. The constituent quark mass is $M = 420$ MeV.

i.e. by rotating the classical saddle point solution $U_c(\mathbf{x})$ through a unitary time-dependent SU(3) collective orientation matrix $R(t)$ and by considering all the collective displacements by $\mathbf{Z}$ of the center of mass of the soliton in coordinate space. The quantization of these collective coordinates will yield the collective wave functions with the appropriate quantum numbers.

The functional integral (12) is thus replaced by a functional integral over R . As described in Ref. [23], we assume adiabatic rotation and small collective velocities in the calculation of the matrix elements for the form factors. This means e.g. that in the expansion of

$$\begin{aligned} D^{-1}\left(R(t)U_c(\mathbf{x} - \mathbf{Z}(t))R^\dagger(t)\right) &= \\ &= e^{i\mathbf{Z}\cdot\mathbf{P}}R\left[D(U_c) + R^\dagger\dot{R} + \gamma_4 R^\dagger m_8 \lambda^8 R\right]^{-1} R^\dagger e^{-i\mathbf{Z}\cdot\mathbf{P}} \end{aligned} \quad (18)$$

only terms up to linear order in the angular velocity $R^\dagger\dot{R}$ and m_8 are retained.

4. A detailed account on how the electromagnetic form factors are derived within this framework is given in Ref. [25]. Here we would like to point out that the only free parameter of the model is the constituent quark mass M . We use in this study the value 420 MeV, which is found from a best fit to many nucleon observables [23]. The other parameters of the model are the current nonstrange quark mass and the cut-off parameter of the proper-time regularization: They are fixed for a given M in the mesonic sector by requiring the physical pion mass and decay constant to be reproduced by the model. The mass of the strange quark is throughout this work set to 180 MeV.

In Fig. 1 we present the results for G_E^0 and G_M^0 . These figures show a general effect which is found in all our calculation, namely the influence of the Yukawa mass μ , governing the asymptotic behavior of the profile function at large r , $\exp(-\mu r)/r$, on the form factors. While the SU(2) soliton incorporates the asymptotic pion behavior $\exp(-m_\pi r)/r$ naturally, the construction of the SU(3) hedgehog by the embedding the SU(2) soliton (15) forces all other pseudo-Goldstone bosons to share the same asymptotic behavior. Therefore, this treatment is phenomenologically unsatisfactory. It is expected that the kaon influence on

	$(G_E^0 + \beta(Q^2, \theta)G_M^0)/(G_M^{p\gamma}/\mu_p)$		$G_E^s + \beta(Q^2, \theta)G_M^s$	
$\beta(Q^2, \theta)$	0.392	0.308 (Model)	0.392	0.308 (Model)
μ	π - K	π - K	π - K	π - K
$M = 420$ MeV	1.433 - 1.695	1.377 - 1.621	0.103 - 0.071	0.101 - 0.067
HAPPEX	$1.527 \pm 0.048 \pm 0.027 \pm 0.011$		$0.025 \pm 0.020 \pm 0.014$	

TABLE I: The combinations $(G_E^0 + \beta(Q^2, \theta)G_M^0)/(G_M^{p\gamma}/\mu_p)$ and $G_E^s + \beta(Q^2, \theta)G_M^s$ for the HAPPEX kinematics $Q^2 = 0.477$ GeV² and $\theta = 12.3^\circ$. The experimental data are taken from HAPPEX [5].

the asymptotics of the meson fields play an important role in the description of the hyperons, even more so if they contain valence strange quarks, as well as in the specific contribution of the strange quark to the properties of the nucleon. With this in mind we tentatively implement a heavier mass asymptotic behavior of the profile function $P(r)$ (16) by adding a diagonal mass term to the Hamiltonian in Eq. (14) and modifying the perturbative treatment of m_s (or m_8) in Eq. (18) by subtracting the same term. We prefer to look at the interval spanned by results of the more consistent pion asymptotics and the phenomenologically driven kaon asymptotics of $P(r)$ as giving an idea of the systematic model uncertainties stemming from the lack of exact treatment of the SU(3) meson asymptotics. Indeed, it is found that the model nucleon electromagnetic form factors do not appreciably change under such procedure, except for the strange electromagnetic form factors and the electric form factor of the neutron.

In Table 1 we present the results for the combination in Eq. (3) of G_E^0 and G_M^0 and for the combination of G_E^s and G_M^s in Eq. (5) as given by HAPPEX [5]. Table 1 further distinguishes between the cases with the HAPPEX collaboration phenomenology based on value $\beta = 0.392$ and the model calculation of the quantity β , as given by

$$\beta(Q^2, \theta) = \frac{\tau G_M^{p\gamma}}{\epsilon G_E^{p\gamma}} \quad (19)$$

(with $\epsilon = [1 + 2(1 + \tau) \tan^2(\theta)]^{-1}$ and $\tau = Q^2/(4M_N^2)$), using the model electromagnetic form factors calculated with the symmetry-conserving quantization. For the combination of G_E^0 and G_M^0 and for both cases of β the HAPPEX result (3) falls within the interval of model results between the pion (π) and kaon (K) asymptotics. The combination of G_E^s and G_M^s seems to overestimate the HAPPEX result (5) for both cases of β . One notices that the kaon asymptotics leads to results closer to the experiment. One has to take into account, however, that the HAPPEX value (5) could be higher, by as much as 0.020, with a different input for the electromagnetic form factors [5]. Although the results in Table 1 depend on M it is found that, for physically reasonable values of M between 400 and 450 MeV the dependence on M is much weaker than the effect of the meson asymptotics.

The results for the strange electric form factor G_E^s are shown in Fig. 2 and for the strange magnetic form factor G_M^s in Fig. 3. For G_E^s the result is basically the same as in Ref. [20] while the G_M^s is strongly modified, compared to [20], to a larger extent by removing a numerical error, as reported in Ref. [26], and to a lesser extent by using in the present investigation the scheme of the symmetry-conserving quantization. Only in the case of G_M^s is a direct comparison with experiment possible. The SAMPLE value (1) [4] is shown in Fig. 3: The model value is not far from the experimental one, but the error bar is too big to draw any other conclusion. The fact that a positive value is obtained, and in particular that




FIG. 2: The strange electric form factor G_E^s as a function of Q^2 . Conventions and model parameter as in Fig. 1.




FIG. 3: The strange magnetic form factor G_M^s , in units of the physical n.m., as a function of Q^2 . Conventions and model parameter as in Fig. 1. The experimental data are taken from SAMPLE [4]

the strange magnetic moment turns out positive, is in agreement with the model independent analysis of [26], which finds $\mu_s = (0.41 \pm 0.18)$ n.m. on the basis of the SU(3) model algebra with input from the hyperon magnetic moments.

In Table 2 we list the χ QSM results for the strange electric and magnetic radii, defined as

$$\langle r^2 \rangle_E^s = -6 \left. \frac{dG_E^s(Q^2)}{dQ^2} \right|_{Q^2=0}, \quad (20)$$

$$\langle r^2 \rangle_M^s = -\frac{6}{\mu_s} \left. \frac{dG_M^s(Q^2)}{dQ^2} \right|_{Q^2=0}, \quad (21)$$

and the strange magnetic moment, for $M = 420$ MeV and both the pion (π) and kaon (K) asymptotics.

μ	π	K
$\langle r^2 \rangle_E^s$ [Fm ²]	-0.220	-0.095
μ_s [n.m.]	0.074	0.115
$\langle r^2 \rangle_M^s$ [Fm ²]	0.303	0.631

TABLE II: The strange magnetic moment and the mean-square strange radii for $M = 420$ MeV.




FIG. 4: The neutral magnetic form factor G_M^s , in units of the physical n.m., as a function of Q^2 . Conventions and model parameter as in Fig. 1. The experimental data are taken from SAMPLE [4, 27].

Using Eq. 2 it is possible to obtain the model prediction for G_M^Z , which is given in Fig. 4, where the experimental result [4]

$$G_M^Z(Q^2 = 0.1) = (1.49 \pm 0.29 \text{ (stat.)} \pm 0.31 \text{ (syst.)}) \text{ n.m.} . \quad (22)$$

is also plotted. Again the χ QSM result falls within the (admittedly large) error bars.

Lastly, Table 3 shows the full model predictions, *i.e.* using both the strange form factors and the electromagnetic ones, for the kinematics of future HAPPEX and A4 experiments. This table is made for the the combination $(G_E^s + \beta G_M^s) / (G_M^{p\gamma} / \mu_p)$ of form factors because it is closer to the quantities extracted from the asymmetries and it can reliably be calculated in the model since this describes well the Q^2 dependence of the form factors [21, 23]. For the sake of consistency we calculate the factor β as given by Eq. (19) using the model. Again we see that the pion asymptotics deviates more from the data than the kaon one, which, nevertheless, seems to overestimate the data. Apart from having to rely on more data to confirm whether this trend will persist, one has to bare in mind that the strange form factors as extracted with the help of Eq. (2) contain a piece due to isospin symmetry breaking [28]. The isospin symmetry breaking effects were considered here to be small on account of large N_c .

5. In this work, we have investigated the strange properties of the nucleon: we calculated $G_{E,M}^0$, the strange form factors $G_{E,M}^s$ and their associated radii, and the strange magnetic moment μ_s within the framework of the SU(3) χ QSM. On the whole, the results of the model

Exp.	A4	HAPPEX II	A4	HAPPEX	A4
$Q^2(\text{GeV}^2)$	0.10	0.11	0.23	0.48	0.48
$\theta(^{\circ})$	35	6	35; 145	12.3	145
β (Mod.)	0.105	0.059	0.255	0.308	0.587
μ	π -K	π - K	π - K	π - K	π - K
$M = 420$ MeV	0.088-0.048	0.091-0.046	0.174-0.108	0.278-0.185	0.297-0.213

TABLE III: The values of $(G_E^s + \beta(Q^2, \theta)G_M^s)/(G_M^{p\gamma}/\mu_p)$ at various Q^2 and θ for $M = 420$ MeV. with pion (π) and kaon (K) tails. All the experiments (Exp.) are still being performed except for HAPPEX.

calculation with kaon asymptotics turn out to be in better agreement with experiment than those with the pion asymptotics, as expected for strange quantities. The results for the kaon asymptotic turn out to be in fairly good agreement with the available data, apart from an apparently slight overestimation of the available HAPPEX data till now on $G_E^s + \beta G_M^s$. The difference between kaon and pion asymptotics gives an indication of the systematic error of the χ QSM in calculating strange quantities. Future data from SAMPLE, HAPPEX, and also from the G0 (JLAB) and A4 (MAMI, Mainz) collaborations, will allow one to judge further the results of the chiral quark-soliton model.

Acknowledgments

The authors are grateful to thank F. Maas (MAMI A4) and M. Pitt (SAMPLE) for clarifying experimental related points and to P. Pobylitsa, M. Polyakov, M. Praszalowicz, and M. Vanderhaeghen for useful discussions and critical comments. AS acknowledges partial financial support from Praxis XXI/BD/15681/98. The work of HCK is supported by the Korean Research Foundation (KRF-2000-015-DP0069). This work has partly been supported by the BMBF, the DFG, the COSY-Project (Jülich), and POCTI/32845/FIS/2000.

-
- [1] D. B. Kaplan and A. Manohar, Nucl. Phys. B310 (1988) 527.
 - [2] M.J. Musolf *et al.*, Phys. Rep. 239 (1994) 1.
 - [3] B. Mueller *et al.* [SAMPLE Collaboration], Phys. Rev. Lett. 78 (1997) 3824.
 - [4] R. Hasty *et al.* [SAMPLE Collaboration], Science 290 (2000) 2117.
 - [5] K. A. Aniol *et al.* [HAPPEX Collaboration], Phys. Lett. B509 (2001) 211.
 - [6] D.E. Groom *et al.* [Particle Data Group], Euro. Phys. J. C15 (2000) 1.
 - [7] R. L. Jaffe, Phys. Lett. B229 (1989) 275.
 - [8] H. Forkel, Phys. Rev. C56 (1997) 510.
 - [9] H.-W. Hammer and M.J. Ramsey-Musolf, Phys. Lett. B416 (1998) 5.
 - [10] N.W. Park and H. Weigel, Nucl. Phys. A541 (1992) 453.
 - [11] H. Forkel, M. Nielsen, X. Jin, and T.D. Cohen, Phys. Rev. C50(1994) 3108.
 - [12] H. Weigel, A. Abada, R. Alkofer, and H. Reinhardt, Phys. Lett. B353 20.
 - [13] S. Hong, B. Park and D. Min, Phys. Lett. B414 (1997) 229.
 - [14] Ulf-G. Meißner, V. Mull, J. Speth, and J.W. Van Orden, Phys. Lett. B408 (1997) 510.
 - [15] T. R. Hemmert, Ulf-G. Meißner, and S. Steininger, Phys. Lett. B437 (1998) 184.

- [16] T.R. Hemmert, B. Kubis, and Ulf-G. Meißner, Phys. Rev. C60 (1999) 045501.
- [17] L. Hannelius, D.O. Riska, and L. Ya. Glozman. Nucl. Phys. A665 (2000) 353.
- [18] N. Mathur *et al.*, Nucl. Phys. Proc. Supl. 94 (2001) 311.
- [19] D. B. Leinweber and A. W. Thomas, Phys. Rev. D62 (2000) 074505.
- [20] H.-Ch. Kim, T. Watabe, and K. Goeke, Nucl. Phys. A616 (1997) 606.
- [21] M. Praszalowicz, T. Watabe, and K. Goeke, Nucl. Phys. A647 (1999) 49.
- [22] D. Diakonov, V. Y. Petrov, and P. V. Pobylitsa, Nucl. Phys. B306 (1988) 809.
- [23] C. V. Christov *et al.*, Prog. Part. Nucl. Phys. 37 (1996) 91.
- [24] R. Alkofer, H. Reinhardt, and H. Weigel Phys. Rept. 265 (1996) 139.
- [25] H.-Ch. Kim, A. Blotz, M.V. Polyakov, and K. Goeke, Phys. Rev. D53 (1996) 4013.
- [26] H.-Ch. Kim, M. Praszalowicz, M. V. Polyakov, and K. Goeke, Phys. Rev. D58 (1998) 114027.
- [27] M. Pitt, a private communication on behalf of the SAMPLE collaboration.
- [28] B.-Q. Ma, Phys. Lett. B408 (1997) 387.